



GIS

جامعة مؤتة، 2014

.....

أ
ب
ج
هـ
ز
ط
ي
ك
ل

:

1	1.1
3	2.1
4	3.1
5	4.1
6	5.1
8	6.1

:

14	2.1
17	2.2
19	3.2
24	4.2
32	5.2
36	6.2

	:
43	1.3
44	2.3
44	3.3
49	4.3
	:
52	1.4
58	2.4
90	3.4
106	4.4
110	2.5
112	
117	

45		1
47		2
47		3
48		4
48		5
49		6
65	(/)	7
65		8
75		9
79		10
83		11
83		12
84		13
91		14
92		15
94		16
96		17
97		18
99		19
101		20

103

21

105

22

7	.	1
8	.	2
53		3
54		4
55		5
56		6
57		7
62		8
62		9
63		10
64		11
67		12
68	(/) (Cd)	13
70	(/) (Cr)	14
71	(/) (Cu)	15
73	(/) (Zn)	16
74	(/) (Ni)	17
76		18
77	(μS/cm)	19
78		20
82		21
84		22
85		23
86		24
87	(ppm) (pb)	25

89	(ppm) (zn)	26
90	(ppm) (Cu)	27
91		28
93		29
94		30
96		31
100		32

27	1
30	2
31	3
31	4
59	5
59	6
60	7
88	8

الملخص

GIS

2013

GIS

GIS

()

.

.

()

.

"

"

"

"

/ (29.35-13.94)

/ (81.09-44.94)

.

Abstract
Environmental Impact of large industrial Plants in the Southern
Region in Jordan Using "GIS"
Nassar Albataiha
Mu'tah University, 2013

The aim of this study is to identify the environmental impacts of large industrial in the southern region of Jordan using Geographic Information System GIS, through a field survey of these facilities, and to show on maps and categorized by activity, and the statement of the spatial scale of the environmental impact on the surrounding areas using GIS, and the aim of this study also, is to identify the health and social problems that plague our population in the surrounding areas and the associated environmental pollution from industrial plants, as well as identify the impact of environmental pollution of the factories on natural resources (soil, water) in the region, so as to get to know the size of these problems and their impact on population.

To achieve the objectives of the study, descriptive analytical method was employed in the analysis and description of spatial data for the sites of industrial installations in the southern region using a number of statistical test provided by geographical information systems to determine the spatial domain of environmental pollution in the region. The study relied on social survey method to collect data from a sample of the study using the study tool (questionnaire), to obtain data on the environmental impacts of industrial plants on the population in the study area.

The result showed that the scope of the impact of environmental pollution of industrial plants "Qatranah Cement factory, national poultry factory, Rashadiyah Cement factory, Alshideah phosphate, and Hasa and Abyed" has covered large parts of the residential areas adjacent to these plants and in particular cities of "Qadisiyah and Rashadiyah, Qatranah and Hasa," and suffer residents of these areas of smoke and thick dust laden with heavy metals, and the result showed that most of the soil and water samples collected from different parts of the study area is contaminated with heavy metals such as lead, Cadmium, Chromium, Zinc, Nickel, Copper, compared with the permissible limits for the concentration of these elements in the soil, Where the concentration of chromium in soil samples (13.94-29.35) mg / kg, and the component lead (44.94-81.09) mg / kg. Also result showed that residents of neighboring areas of industrial plants in the study area are suffering from the spread of diseases such as allergic skin and eyes, and sinus diseases, respiratory diseases, diseases of tonsillitis and chronic ear.

1.1

:

1992

"

"

1972

2002

.(2003)

(ETA)

.

.

1995

1980

.

1996

.

2003

2003 (1)

.2006 (52)

.(2006)

$$\frac{60}{1.66}$$

$$\frac{()}{()}$$

:

:

-1
-2
-3

: **3.1**

:

(1

GIS

(2

(3

()

(4

(5

()

(6

(7

4.1

الصناعات

3

التعدينية والكيميائية

5.1

-1

-2

(1)

:

% 50

% 9.2

502000

(2012

)

:


-3

(2)


-4

% 10

-5



(1)



(2)

6.1

.

.% 70

%30

.()

-1

36 14 17 35 19 53

31 28 31 30 53 03

(%3.9)

2 3504

.

:

(150 -50)

()

350

(200-100)

.

[35 18 03 – 36 05 55]

(1) [30 36 16 – 31 0 07]

2 2209,08

.

-

-

.

.

. 222 (2013)

1150 6.5

3 -
12-10

. 23.6

：

(132) (360) .

)

.(2013

3 (8)

.(2005) 3 (14)

(30 30 31 27)

(35 32 35 27)

(210)

(%36.98) ² (32832)

.

()

()

：

： .1

.(%95)

： .2

(150)

.) (2009)

28

36

4

2

.

-

300

(250 -200)

500

)

(100)

.) (2009

.

1.2

(Koop & Tole, 2004)


(Ostro, 2006)

)

(2000

-(1999)

.	Output	Input	- 1
.	Output	Product	-2



\cdot
 -2
 \cdot
 -3
 \cdot
 (NO_x) :
 $($) (SO_4)

\cdot
 -4

\cdot
 -5

\cdot
 $)$
 $($

)
(

-6

.

-7

.

(2000)

2.2

)

.(2013

:

.

:

(1

:

(2

.

-

-

.

.

.

(164) (84)
 (70) 2005
 (2012) (149)

.

.

3.2

)

(
)

.(2002

.(()

)

.(2002

250

%5

.(2013)

:

-1

.(2006)

(2009)

)

.(2000

.(2002)

(1948)

(17)

(2009)

Suspended Particulates (SP) -

-:

Oxides of Carbon (Cox) .1

Oxides of Nitrogen (Nox) .2

Sulphur Oxides (Sox) .3

Hydrocarbons .4

. -2

%98 %71

%97 .

%3

(1999)

.(1990)

)

(

.(2002)

:

(1

(2

(3

(4

(5

4.2

.

)

(2009

.

)

(2009

(Merenu, 2003)

2006

5

90

5

300

14001

.(2014)

"

"

.

.

.

.

()

1951

300 2011

% 98

(2013)

—

:

—

—

(1)



(1)

2011 7.6

1953

1.864 882

150

55

2008 . 4.847

% 48

) 2012 7.6

.(2012

(%72-63)

.(2005)

.

()

10

.

2013

%3.5

%4

(35 - 20)

.

.

.

6 %20

.

.()

.

(4) (3) (2)



(2)



(3)



(4)

.(2013)

5.2

1964 (GIS)

.(2009)

.

.

(Levick et al. 2000)

:

.(Thomas et al, 2003)

.(Azam, 2002)

.(2002)

.

.

()

.(United Nations, 2006)

.(2008)

.

:

:

.

:

.

:

.(Aurom, 1999)

:

:

.

:

()

.(Ray, 2000)

.

.

Industrial "

Industrial "

"Cluster Concentration"

"Complex

. "Agglomeration

.

.(1995)

.

.(1985)

.(2005)

: **6.2**

:

" (2005)
" GIS

.

" (2000)
"

.

Arc GIS 9.3

% 5

.

% 44 % 41

.

" (2006)
"

(DPSIR)

(Driving Force)

(Impacts)

(State)

(Pressure)

.(Response)

·
(Dredging)

·
· 2.5

()
%7

·

·
" (1999)
"

·

·
%10

·
:
(1
(2

·
(3

·
(4

·
(5

·
" (2010)

"

.1

)

(

.

.2

.

.3

.

.4

.

.

"

(2004)

"

.

.

.

.

•

•

11

•

•

(88)

•

.1

•

(

•

.2

•

.3

1.3

:

" "

.

.

.

ArcGIS 10

.Buffer Zone

GPS

.

GPS

2.3

:

()

:

-1

:

100000 :1

-2

Google Earth Pro

-3

-4

:

:

()

(1

(2

3.3

)

(2360)

(

(1)

% 18.50

437

. 2013

. "

"

(1)

(1)

الرقم	عدد الأسر الكلي	الأهمية النسبية	%
1	880	127	14.43
2	530	95	17.92
3	470	97	20.64
4	480	118	24.58
	2360	437	18.52

:

:

.

:

:

:

:

:

.

:

.

:

4

:

.

(4)

.

:

.

-

(2)

(%)	
7.50	33
40.40	177
24.10	105
28.00	122
100.00	437

(2)

%40.4

%24.1

%28.0

.%7.5

-

(3)

(%)		
14.50	63	24-18
33.40	146	35 -25
34.10	149	45-36
18.00	79	55 -45
100.00	437	

(3)

-25)

%34.1

(45-36)

% 33.4 (35
%18.1 (55-45)
.%14.50 (24-18)

-

(4)

(%)	
74.6	326
25.4	111
100.00	437

(4)

.%25.4 %74.6

-

(5)

(%)	
10.10	44
41.80	183
30.90	135
8.80	38
8.40	37
100	437

(5)

%41.8

% 30.9

% 8.8

%10.1

.%8.40

-

(6)

(%)		
11.80	52	4
35.90	157	6-4
33.90	148	8-6
18.40	80	8
100	437	

(6)

(8-6)
)

%35.90

(6-4)

% 33.9

%18.40

(8

.% 11.8

(4)

4.3

:

:

100000 :1

Arc

Google Earth Pro.

GPS

GIS 10

.

:

" "

.

:

SPSS

.

-1

-2

:

()

:

.

.

120

20

85

.

.

:

(EC)

(1)

6

(5)

(conductivity meter Ec-meter model LF, 750)

(pH)

(5) (1) (pH)

6

.(pH meter model HI metet)

(CaCO₃)

.(Bashour, 2007)

(Total Organic Matter)

.Walkley- Blacks

:
)

(

.

.

1.4

Buffer Distance

Buffer Distance

()

(3)

(5)

(5-3)

:


"

"

Multiple Buffering

(3) (4).


1



(3)

Google Earth Pro.

:



(4)

Google Erath Pro.

:

(3)

10

:


(3)

(3)

12

(4)

(5)



(5)


Google Erath Pro.

:

Multiple Buffering

1

(6).



(6)

Google Erath Pro.

:

(6)


(6)

(3-0)

Multiple Buffering

6 1

.(7)



(7)

Google Erath Pro.

:

(7)

(7)

(5-3)

.(5)

2.4

:

:

.

.

()

.


)

(

(6) (5)

.


.



(5)



(6)



(7)

. :

:

.

:

:

-1

:

.

: -2

:

.

:

(Organic Contaminants) .1

.

Inorganic Contaminants .2

.


.

.

.

(11) (10) (9) (8)

.



(8)

Google Earth Pro


:



(9)

Google Earth Pro


:



(10)

Google Earth Pro

:



(11)

Google Earth Pro

:

.

() .

As,Sr,Pb,Hg, Cd, Ni, Se, Cr Al, F, Cl :

.

:

3 5

38

:

.(Azam,2002)()

(8)

(7)

جدول (7)

تركيز العناصر الثقيلة في عينات التربة (ملغرام/ كغم)

الحديد Fe	الرصاص Pb	النيكل Ni	الزنك Zn	النحاس Cu	الكروم Cr	الكاديوم Cd	العينة
51.2	67.15	6.85	68.3	3.5	26.31	29.1	1
52.3	65.09	7.22	66.28	3.8	28.31	27.42	2
45.67	79.62	8.2	65.94	2.4	25.61	24.53	1
43.54	77.43	7.23	61.37	2.9	27.61	21.54	
33.7	80.08	8.29	62.37	2.5	29.35	27.38	1
39.76	81.09	6.72	66.92	3.3	31.28	23.94	2
31.26	56.23	7.6	70.26	3.9	20.11	0.3	1
30.25	47.84	7.74	70.45	4.1	19.3	0.1	2
32.56	45.32	7.88	71.11	4.6	18.66	0.2	3
55.83	50.12	8.2	68.36	3.9	15.6	0.4	1
54.75	46.35	8.5	69.11	4.3	14.82	0.3	2
55.11	44.94	8.4	70.12	4.5	13.94	0.1	3

(8)

(mg/kg)
500
500
16.66-3.33
1.66
3.33
8.33
8.33 - 16.66

المرجع: (UNEP, 1991)


.

.

. (Koop & Tole, 2004)

-44.94

(4) / 81.09



(12)

المصدر: عمل الباحث.

:

-1

-2

-3


Cadmium (Cd)

()

.(Ostro, 2006)

النتائج:

(13) (mg/kg) 29.1 -0.1



شكل (13)

تركيز عنصر الكاديوم (Cd) في عينات التربة بالملغرام / كغم

المصدر: عمل الباحث.

:

-1

-2

-3

-4

(Cr) Chromium


:

:

.(Singh, 2005)

29.35 -13.94

.(14) /



شكل (14)

تركيز عنصر (Cr) في عينات التربة بالملغرام / كجم

المصدر: عمل الباحث.

:

-1

-2

(Cu) Copper

:


:

(Wilson)

.(Al-Momani, 2003)

:

.(15) / 4.9 -2.4



شكل (15)

تركيز عنصر النحاس (Cu) في عينات التربة بالملغرام / كغم

المصدر: عمل الباحث.

:

-1

-2

.

(ZN) Zinc


:

. (Azam, 2002)

.

:

.(16) (mg/kg) 71.11 -61.37



شكل (16)

تركيز عنصر الزنك (Zn) في عينات التربة بالملغرام / كغم

المصدر: عمل الباحث.


(16)

(Merenu, 2003)

:

:

(17) (mg/kg) 8.5 - 6.72



شكل (17)

تركيز عنصر النيكل (Ni) في عينات التربة (الملغرام / كغم)

المصدر: عمل الباحث.

(17)

(9).

(9)

العينة	درجة الحموضة (pH)	الموصلية الكهربائية ($\mu\text{S/cm}$)	مجموع المادة العضوية %
1	8.23	557.62	11.86
2	7.92	562.3	11.35
3	7.46	542.9	12.31
1	9.01	659	2.35
2	8.35	519.66	2.69
1	8.02	328	4.21
2	8.29	398	3.89
1	9.14	223	3.23
2	8.31	344	3.68
1	8.36	434	2.31
2	8.42	533	1.98
3	8.55	239	2.6

المصدر: عمل الباحث.


(pH)

(1)

8.29 (9.14 - 7.46)

(18)

(1)



(18)

المصدر: عمل الباحث.

(Momani,2003)

.(Merenu ,2003)

:(E.C)

)


(ms/m)

:

(2004

($\mu\text{S}/\text{cm}$)

(19).



(19)

($\mu\text{S}/\text{cm}$)

المصدر: عمل الباحث.

(19)

($\mu\text{S}/\text{cm}$) 444.8

/

659

223.0

(1)

(19)


(1)

:

(Ostro, 2006)

(20)

(9)



(20)

المصدر: عمل الباحث.

(20)

% 4.99

% 12.31-1.98

(2)

% 80 % 5

.(Thomas et al. ,2003)

% 73 - % 30.4

.(2)

جدول (10)
نسبه كربونات الكالسيوم في عينات التربة

(%)	
47	1
37.5	2
31.5	3
36.5	1
30.5	2
32	1
38.5	2
42	1
40	2
70	1
68	2
30.4	3

المصدر: عمل الباحث.

:

296

%71

% 98

% 97

%3

.

.

.

)

(

.

:

(1

.

(2

.

(3

.

(4

.

—

—

.

.

.


.

.

:

(21)

.



(21)

المصدر: عمل الباحث، بالاعتماد على برنامج Google Earth Pro.

(11) (12)

(13)

(11)

المحتوى العضوي %	الموصلية الكهربائية ($\mu\text{S}/\text{cm}$)	الحموضة (pH)	العينة
7.52	835	8.29	-
12.33	1069	8.64	1
13.61	1704	7.9	2
14.38	1616	8.36	
11.62	511	8.39	1
11.95	569	8.22	2
13.91	1251	8.15	
10.37	907	7.95	x1
6.62	885	8.1	1
6.73	893	7.86	2

(12)

النحاس Cu ppm	الحديد Fe ppm	الزنك Zn ppm	الرصاص Pb Ppm	العينة
2.4	0.75	0.17	0.26	-
1.8	0.72	0.15	0.18	1
1.6	0.68	0.13	0.16	2
2.2	0.72	0.15	0.27	
0.8	0.65	0.25	0.25	1
3.2	0.62	0.23	0.29	2
2.4	0.81	0.19	0.32	
2.2	0.76	0.22	0.24	x1
2.8	0.62	0.18	0.26	1
2.2	0.63	0.21	0.27	2


(13)

(ppm)
0.01
5
1

المرجع: (UNEP, 1991)

(pH) :

(22) 8.64 7.86
(2) (1)



(22)

.


)
(5)

(2002

(23) (μS/cm) 1704 511

(2)

.(1)



(23)

: المحتوى العضوي

(6)


% 14.38-6.62

(19)

(1)

)

.(2008




(24)

(ppm) 0.32 - 0.16

(2)

(ppm) 0.01



(25)

تركيز عنصر الرصاص (ppb) لعينات المياه

:

(1

(2

(3

(8)



(8)

:


:

(ppm) 0.25 - 0.13

(2)

(1)

(ppm) 5



(26)

تركيز عنصر الزنك (zn) (ppm) لعينات المياه

:

:

:

(1)

(2)

(3)


(4)

:

(ppm) 3.2 - 0.8

(2)

(ppm) 1



(27)

تركيز عنصر النحاس (Cu) (ppm) لعينات المياه

:

3.4


-

(28)

(14)

(14)

%	
49.2	215
9.5	42
16.1	70
20.9	91
4.3	19
100	437



(28)

(28)


(14)

% 49.2

% 20.9
 % 16.1
 " % 9.5
 .%4.3 "
 % 54.77
 41.29
 %
 .
 (29) (15)
 .
 (15)

%	
15.1	66
28.5	125
34.1	149
22.3	97
100	437

. :



(29)

(29)

(15)

% 34.1

% 28.5

%22.3


% 15.1

(30)

(16)

(16)

%		
16.9	74	()
39.4	172	
15.8	69	
19.5	85	
8.5	37	



(30)

(30)

(15)

% 39.4

% 19.5

()

% 16.9

.%8.5

%15.8

(17)

% 19.5

% 8.5


.

.

.

(17)

%		%		%		%		%	
2.5	11	3.9	17	3.4	15	9.8	43	9.4	41
1.8	8	5.7	25	4.1	18	2.7	12	7.3	32
1.6	7	3.7	16	3.4	15	2.3	10	11.2	49
2.5	11	6.2	27	4.8	21	2.1	9	11.4	50
8.5	37	19.5	85	15.8	69	16.9	74	39.4	172



(31)

)

(

.

(18)

1	0.874	4.043	1
2	0.788	4.017	8
3	0.863	3.942	7
4	0.857	3.825	6
5	0.981	3.808	5
6	1.031	3.750	4
7	1.024	3.667	3
8	1.296	3.367	2
-	0.687	3.833	1-8

0.687 (3.833) (18)
 " (1) (4.04)
 " (2) (3.367)
 (18) "

(19)

(% 27.9)

% 5

(2013) % 7.4

% 11.7

% 10.7

% 9.3 " "


% 8.3

. % 0.7

(19)

()									
%		%		%		%		%	
1.7	1	16.7	10	15.0	9	10.0	6	13.3	8
1.7	1	13.3	8	15.0	9	13.3	8	10.0	6
0.0	0	15.0	9	13.3	8	10.0	6	11.7	7
0.0	0	13.3	8	10.0	6	8.3	5	11.7	7
0.7	2	11.7	35	10.7	32	8.3	25	9.3	28

. :



(32)

)

(

(20)

1	0.881	4.117	1
2	0.930	4.025	5
3	0.914	3.933	3
4	1.043	3.825	2
5	0.78	3.819	4
-	0.545	3.976	5-1

(20)

0.545

(3.97)

(1)

(4.117)

"

(4)

(3.819)

"

"

(20)

."

.

-

.

(21)

الآثار الاقتصادية للتلوث البيئي

الآثار الاقتصادية للتلوث البيئي			
الآثار الاقتصادية للتلوث البيئي			
1	0.912	4.342	1
2	0.823	4.109	2
3	0.901	3.984	5
4	0.895	3.765	7
5	1.024	3.618	9
6	1.076	3.584	10
7	1.069	3.348	3
8	1.353	3.011	4
9	0.903	3.001	8
10	1.034	2.987	6
-	0.876	3.578	1-10
الآثار الاقتصادية للتلوث البيئي			البيئي

(21)

0.876

(4.342)

"

(1)

(4.342)

"

"

(6)

(2.987)

"

(21)

الإجراءات الحكومية اتجاه الوقاية من التلوث البيئي الصادر من

-

(22)

الإجراءات الحكومية اتجاه الوقاية من التلوث البيئي

الإجراءات الحكومية اتجاه

الوقاية من التلوث البيئي

1	0.862	3.765	7
2	0.766	3.481	10
3	0.850	3.431	3
4	0.844	3.409	6
5	0.984	3.209	8
6	1.041	3.154	5
7	1.033	3.089	4
8	1.142	2.729	9
9	0.853	2.439	1
10	0.995	2.364	2
-	0.823	3.107	1-10
الآثار الاقتصادية للتلوث البيئي			

(22)

0.823

(3.107)

"

(7)

(3.765)

(2)

(2.364)

"

"
.

(22)

.

4.4

:

.

.

()

(5.3-3)

(1

10

(2

(3

% 4.99 % 12.31-1.98

% 14.38-6.62

.

(4

(3.97)

% 11.7

% 10.7

% 9.3

" "

% 8.3

.

% 0.7

(5

0.876

(4.342)

.

(6

0.823

(3.107)

.

5.4

:

(1

.

(2

.

(3

.

(4

.

(5

.

(6

.

-

(2006)

161-120 (117) (31)

.
(2010)

"

"

. 2010 31-30
(2009)

. 78-51
(2009)

. 218-191 31 94
(2003)

.
(1995)

40

.161-160 . 1995
(2008)

.

(2002)

(2000)

(1999)

(2004)

187-163 (2) -(20)

(2009)

(2012)

(1996)

:

. 89 78 1 23

(1999)

(2000)

(2004)

(2009)

(1985)

.

(2010)

. 195-174 (5) (3)
(2000)

.

(2002)

.

() (1990)

.

(2006)

(2) (6)

.

.281 -205

(2002)

.

(2000)

.

(2000)

.

(1999)

. 167-131 (2) (14)

(2005)

.(2008)

(2014)

(2013)

(2006)

(2012)

(2004)

(2002)

(2000)

(2008)

2007-202

(2005)

(2005)

(2013)

- : -
- Al-Momani, I.F., (2003), Trace elements in atmosphere precipitation at northern Jordan measured by ICP-MS :acidity and possible sources, **Atmos Environ.**, 37, 4507-4515.
- Azam. Elhag (2002). **The Analysis of chromium and Hydrogen Sulphide discharged from the tanneries and their effect on the environment**, unpublished M.sc, thesis, Department of Applied Chemistry and Chemical technology, Faculty of Science and technology, U.O.G.
- Aurom, Daivd.(1999) **Regional Economic and analysis for practitioners; An Introduction, to Common Descriptive Methods.** First printing by Prager Publisher. U.S.A.
- Merenu I.A. (2003). **The Effect of Cement Dust Exposure on Lung Function in Cement Factory Workers In Sokoto Metropolis**, unpublished MPH dissertation, Usman Danfodio University, Sokoto, Nigeria.
- Murray, G.Y., Hendershot W.H., (2000), Trace metals speciation and bioavailability in urban soils **Environ. pollution**, 107, 137-144.
- Ray. M. Northman (1975) **Urban Geography**, Origin State University. New York. P 304-311.
- Ostro, B. (2006) "Air Pollution and Mortality, *Journal of Exposure Analysis and Environmental Epidemiology*, Vol. 6, No.1., (88-133)..
- Koop, G., and Tole, L, (2004) "Measuring the health effects of air pollution: to what extent can we really say that people are dying from bad air", **Journal of environmental economics and management**, vol. 47, No.6 (30-54)..
- Levick, L. R. ,Kidwell, M. R., and H. D. Fox (2000) **Integration of Watershed and Ecological Site Land Classification Scales Using GIS Technology**, (ASCE)
- Symeon Christodoulou, Pooyan Aslani, and Annie Vanrenterghem (2003) **A Risk Analysis Framework for Evaluating Structural Degradation of Water Mains in Urban Settings, Using Neurofuzzy Systems and Statistical Modeling Techniques**, ASCE, World Water & Environmental Resources Congress 2003.
- Thomas H. Cahill, P.E., John Hoekstra, Wesley Horner, Susan McDaniels, and Michele Adams, P.E (2003) **Application of a Water Balance Model and GIS for Sustainable Watershed Management**, American Society of Civil Engineers (ASCE).
- UNEP (1991). **Monitoring Industrial Emissions and wastes**, United Nations Publications.
- United Nations (2006) **Industrial Estates in Asia and the Far East** , New York , USA.

()

- () -
 : -
 -11

(:
 : -12

			()

1. الأثر البيئي على صحة السكان

-(/)

						1
						2
						3
						4

2.

أوافق بشدة	أوافق	محايد	لا أوافق	لا أوافق بشدة	الملوثات	
						1
						2
						3
						4
						5
						6
						7
						8
						9
						10

3. الآثار الاقتصادية للتلوث البيئي

أوافق بشدة	أوافق	محايد	لا أوافق	لا أوافق بشدة	الملوثات	
						1
						2
						3
						4
						5
						6
						7
						8
						9
						10
						11

4. الإجراءات الحكومية اتجاه التلوث البيئي الصادر من المصانع

أوافق بشدة	أوافق	محايد	لا أوافق	لا أوافق بشدة	الملوثات	
						1
						2
						3
						4
						5
						6
						7
						8
						9
						10